


Recent Results on Doubly Charmed Baryons

CIPANP May 20, 2003
Peter S. Cooper
Fermi National Accelerator Laboratory

Outline

- > Brief theory and model review
- Search methods and Previous results
- ➤ Where are the isospin partners?
- ➤ New States A pair of Isodoublets?
- ➤ What are these? Properties and production
- **Conclusions**

SELEX Collaboration





VOLUME 89, NUMBER 11

PHYSICAL REVIEW LETTERS

9 SEPTEMBER 2002

First Observation of the Doubly Charmed Baryon \(\mathbb{E}_{cc}^{+} \)

M. Mattson, G. Alkhazov, II. A. G. Atamantchouk, III. M. Y. Balatz, S. A. N. F. Bondar, II. P. S. Cooper, S. L. J. Dauwe, I. G. V. Davidenko, S. U. Dersch, S. A. G. Dolgolenko, G. B. Dzyubenko, R. Edelstein, S. L. Emediato, IS. A. M. F. Endler, J. Engelfried, J. E. Echnich, S. C. O. Escobar, IS. A. V. Evdokimov, S. I. S. Filimonov, IS. F. Emediato, IS. M. N. F. Endler, J. Englefried, J. H. Cooper, J. C. Escobar, IS. A. V. Evdokimov, S. I. S. Filimonov, IS. F. G. Garcia, S. M. M. Gaspero, IS. I. Giller, IS. V. L. Golovisov, IP. Gouffon, IS. E. Gulmez, He Kangling, M. Itori, IS. Y. Jun, M. Kaya, IS. J. Kilmer, V. T. Kim, II. L. M. Kochenda, II. I. Konorov, A. P. Kozhevnikov, A. G. Krivshich, II. H. Krüger, M. A. Kubantsev, V. P. Kubarovsky, A. I. Kulyavisev, S. P. Kucpatkin, IS. I. V. F. Kutshetsov, A. Kushnitenko, S. Kwan, J. Lach, V. P. Kubarovsky, A. I. Kulyavisev, S. D. Kucpatkin, IS. I. Whatham, M. Luksy, I-I. Lungov, IS. V. P. Maleev, II. D. Mao, Sh. Mao Chensheng, Mao Zhenlin, P. P. Mathew, St. V. Marveev, E. McCliment, M. A. Moinester, I. V. Wolchanov, A. Morelos, IS. K. D. Nelson, IS. A. V. Nemitkin, IS. P. N. V. Neoustroev, II. C. Newsom, IS. A. P. Nilov, S. N. Nurshev, A. Ocherashvili, I-Iz. E. Oliveira, Y. Onel, IS. E. Ozel, IS. S. Ozkorucuklu, IS. A. Penzo, O'S. V. Petennko, P. Pogodin, IS. M. Procano, St. V. A. Prutskoi, E. Ramberg, G. F. Rappazzo, B. V. Razmyslovich, I. M. Putzo, O'S. V. Petennko, P. Schiavon, S. J. Simon, S. A. Simikov, D. Skow, V. J. Smith, IS. M. Srivastava, IS. V. Steiner, I. V. Stepanov, II. M. L. Stutte, M. Svoiski, II. N. N. K. Terentyev, S. II. G. P. Thomas, I. L. N. Uvarov, II. A. N. Vasiltev, D. V. Vavilov, V. S. Verebryusov, V. A. Victovsko, V. S. Vishnyakov, A. A. Vorobyov, II. K. Vorwalter, S. J. You, A. Zhao Wenheng, J. Zheng Shuchen, and R. Zukanovich-Funchallon.

(SELEX Collaboration)

¹Ball State University, Muncie, Indiana 47306 ²Bogazici University, Bebek 80815 Istanbul, Turkey ³Carnegie-Mellon University, Pittsburgh, Pennsylvania 15213 ⁴Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil ⁵Fermi National Accelerator Laboratory, Batavia, Illinois 60510 ⁶Institute for High Energy Physics, Protvino, Russia Institute of High Energy Physics, Beijing, People's Republic of China SInstitute of Theoretical and Experimental Physics, Moscow, Russia ⁹Max-Planck-Institut für Kernphysik, 69117 Heidelberg, Germany ¹⁰Moscow State University, Moscow, Russia 11 Petersburg Nuclear Physics Institute, St. Petersburg, Russia 12 Tel Aviv University, 69978 Ramat Aviv, Israel 13 Universidad Autónoma de San Luis Potosi, San Luis Potosi, Mexico ¹⁴Universidade Federal da Paraíba, Paraíba, Brazil 15 University of Bristol, Bristol BS8 1TL, United Kingdom ¹⁶University of Iowa, Iowa City, Iowa 52242 17 University of Michigan-Flint, Flint, Michigan 48502 18 University of Rome "La Sapienza" and INFN, Rome, Italy 19 University of São Paulo, São Paulo, Brazil 20 University of Trieste and INFN, Trieste, Italy (Received 22 July 2002; published 20 August 2002)

We observe a signal for the doubly charmed baryon Ξ_r^+ in the charged decay mode $\Xi_r^+ \to \Lambda^+ K^- \pi^+$ in data from SELEX, the charm hadroproduction experiment at Fermilab. We observe an excess of 15.9 events over an expected background of 6.1 ± 0.5 events, a statistical significance of 6.3σ . The observed mass of this state is $3519 \pm 1 \text{ MeV}/c^2$. The Gaussian mass width of this state is $3 \text{ MeV}/c^2$, consistent with resolution; its lifetime is less than 335 at 79% confidence.

DOI: 10.1103/PhysRevLett.89.112001


PACS numbers: 14.20.Lq, 13.30.Eg


Some Nomenclature


SELEX

Searching for charm laryous since 1995

In this talk we replace PDG names with suggestive labels. While SU_4 is badly broken by the charm mass it still classifies all the states .There are many model predictions in the 3.5-3.8 GeV range







Author	Year	Model	$\Xi cc(J=3/2)$	$\Xi cc(J=1/2)$
Bjorken	1986	Phenom	3.70 GeV	3.64 GeV
Fleck & Richard	1989	Bag	3.636	3.516
Fleck & Richard	1989	Quarkonium	3.741	3.613
Roncaglia	1995	Feynman/Hellamn	3.81	3.66
Ellis	2002	Phenom	3.711	3.651

Sampling of Models


Overall Features


- ground states near 3.6 GeV
- ground states I=1/2 multiplets degenerate
- Hyperfine splitting around 60-120 MeV
- Most models predict on electromagnetic hyperfine transitions
- Some models predict pionic transitions for J=3/2 J=1/2
- Model dependent predictions for orbital and radial excitations

SELEX Experiment at Fermilab

SELEX SELEX Searching for charm buryons since 1905 SELEX Searching for charm buryons since 1905 SELEX SELEX

Data taken 1996-7 in P-Center @ FNAL





SELEX (E781)

SELEX Experiment


- > Forward charm production $x_F > 0.1$
- \rightarrow π^{-} p and Σ^{-} beams @ 600 GeV
- > Typical boost ~100

+0.76


- > RICH PID above 22 GeV
- 20 plane 4 view svx σ>4 μm

Peter S. Cooper - Fermilab

Experimental Evidence from 2002




SELEX reported 3 significant high mass peaks



We argued that these states are doubly-charmed baryons


Search strategy and results



- ightharpoonup ccq weak decays into csq. Look for charm, strange and baryon in the final state, starting with Selex's L_c^+ sample: L_c^+ $K^ p^+$, L_c^+ $K^ p^+$ p^+
- ► Look for new secondary vertex between primary and L_c⁺
- ➤ No RICH PID on new K pt tracks (5-20 GeV/c too soft)
- All cuts set (and fixed!) from previous searches (e.g. $L/\sigma > 1$)

ccd⁺(3520) results

- Fix L_c^+ mass to 2284.9 MeV/c² (PDG)
- Right sign has a mass peak at 3520 consistent with resolution. Wrong sign ($L_c^+K^+p$) has no structure.
- >15.8 signal, 6.2 background, $15.8/\sqrt{6.2} = 6.3$ s
- ➤ Possion Prob=1x10⁻⁶; anywhere, 1.1x10⁻⁴



ccd⁺(3520) Iso-partner?


- \triangleright Same cuts as ccd⁺ ~3.5 σ hint in L_c^+ K⁻ p⁺ p⁺
- ➤ Nothing in wrong sign: L_c+K+p-p*
- > cos θ_K^* for mass sidebands (soft vtx tracks). Set cut with $S_{MC}/\text{sqrt}(B_{tot})$, (cos $\theta_K^*>-0.6$)
- \triangleright MC signal is flat, expect: $s \rightarrow 0.8s$, $b \rightarrow b/3$


ccu⁺⁺results


- ➤ Right sign has a mass peak at 3460 consistent with resolution, L=0. Wrong sign ($L_c^+K^+pp$) no structure.
- \geq 7.1 signal, 0.9 background, 7.5 σ , Possion Prob <10⁻⁵
- > 60 MeV is **too big** for an isospin splitting.


Doubly charmed baryons: $\Xi_{cc}^{++}(3460)$, $\Xi_{cc}^{+}(3520)$

Hard to understand these as an Isodoublet










Where are the Iso-partners?


- \triangleright MC signal for phase space $\operatorname{ccd}^+(3520)$ decay is flat
- \triangleright What happens if we apply $\cos \theta^*_{K} > 0.6$ cut here?
- $ightharpoonup \operatorname{ccd}^+(3520)$ strongly attenuated: 16 / 6 \circ 5 / 1
- $ightharpoonup \operatorname{ccd}^+(3520)$ clearly not like phase space \mathbb{R} L>0
- ➤ Not Isodoublet with 3460 different angular dist


New ccd⁺(3443) candidate now very significant

- \triangleright Before cut we ignored bump at 3443 only 4σ
- ➤ Background departs bump at 3443 remains
- \triangleright 7.4 signal, 1.6 background, 5.8 σ , Prob<3.8 \times 10⁻⁵
- ightharpoonup Consistent with L = 0: 11 / 7 ® 7.4 / 1.6
- \triangleright A partner for the ccu⁺⁺(3460) \triangle M=17 MeV **CIPANP - 2003** Peter S. Cooper - Fermilab











ccd⁺(3520) Iso-partner?


- $ightharpoonup \operatorname{ccd}^+(3520)$ mostly has $\cos \theta^*_{K} < -0.6$
- $ightharpoonup L_c^+$ and $m K^-$ back-to-back: $\cos \theta^*_{K} \cos \theta^*_{\Lambda c} < -0.25$ keeps 90% of the $\rm ccd^+(3520)$ signal
- ➤ Apply to the ccu⁺⁺ sample to search for an Isopartner for the ccd⁺(3520)
- No signal / little background with $L/\sigma > 1$. Try reducing cut to $L/\sigma > 0.25$

New ccu⁺⁺(3443) candidate now very significant


- > Bump at 3443 remains background departs
- \geq 7.4 signal, 1.6 background, 5.8 σ , Prob<3.8 \times 10⁻⁵
- \triangleright Consistent with L > 0
- ightharpoonup A partner for the ccd+(3520) Δ M=21 MeV CIPANP 2003 Peter S. Cooper Fermilab










Recap - A Pair of Isodoublets?










CIPANP - 2003

Peter S. Cooper - Fermilab

Lifetimes



> Selex uses reduced proper lifetime technique


$$c\mathbf{t} = m/p_z[L-L_{\min}]; L_{\min} = \mathbf{s}$$

- ➤ Make simulation templates for different lifetimes
- > Lifetimes of all states near our resolution limits

<30 fsec; 0 not ruled out

Model Predictions

- ➤ Guberina, et.al. HQET + 1/M_O
 - $\tau[\Xi_{cc}^{+}] \sim 200 \text{ fsec}$
 - $\tau[\Xi_{cc}^{++}] \sim 1000 \text{ fsec}$
- Observed states don't seem to follow predictions
- ➤ How can the decay rate for ccq states be so large?



		1			4		
P	rn		11	P1			n
	LU	u	u		LΙ	v	

	_		Searching for charm baryons since 1996 literaty of but thereby of factors:
Beam Hadrons	S-	proton	b.
Luminosity fraction	0.77	0.13	0.10
ccu ⁺⁺ (3443) signal/ sidebands	6 /10	2/2	0 / 1
ccu ⁺⁺ (3460) signal/ sidebands	8 / 9	3/0	0 / 0
ccd ⁺ (3520) signal /sidebands	18/18	4 / 1	0 / 1
ccd ⁺ (3541) signal /sidebands	7/10	4 / 1	0 / 1
	86	17	3 (11?)

Dominantly produced by baryon beams

for ccd+ $\langle x_p \rangle \sim 0.35$ (200GeV), $\langle p_t \rangle \sim 1$ GeV/c - like single charm SELEX

- > Focus (E831) has looked in 250 GeV/c photo-production NO signals with 20K L_c^+ , many other modes searched no signals anywhere
- > E791 has looked in 250 GeV/c pp production no signals

Conclusions



- > Selex reports 4 high mass baryon states in an apparent pair of Isodoublets.
 - Lower mass doublet is consisent with L = 0 decay
 - Upper mass doublet is inconsistent with L = 0 decay
- > All 4 states decay like doubly charmed baryons with very short lifetimes (<30fs)
- ➤ The splitting (~19 MeV) is large for an isodoublet.
- \triangleright Radiative decays (e.g. 3520 ® 3443 γ) suppressed below weak decay rate?


Where do these states fit into our theoretical framework?

➤ Bardeen, Eichten and Hill and suggest these might be the spectroscopy of the cc "nucleus" of a ccq baryon:

$$J^{P} = 1/2^{+} \left[\mathbf{c} \uparrow \mathbf{c} \uparrow L = 0, J^{P} = 1^{+} \right] q \downarrow \qquad \qquad J^{P} = 1/2^{-} \left[\mathbf{c} \uparrow \mathbf{c} \downarrow L = 1, J^{P} = 1^{-} \right] q \downarrow$$

- ➤ Predicted splitting consistent with observed 78 MeV
- First EM transition is M2. Decay rate ~ 1/1.5 [fsec]

Conclusions (2)




Other Interpretations


- > Narrow very high mass singly charmed baryon states?
 - Like Babar's D_S*(2317) and Cleo's D_S*(2460) ?
 - Quark content of $L_c^+K^-p^+p^+$ is a pentaquark $\left[csu\overline{d}u\right]$ not a baryon
 - Why should there be ultra narrow baryon states with Q~500 MeV?

Open questions

- **Production**
 - 1/2 of Selex's L_c^+ of come though these states
 - Only seen in forward production by baryons, no π or γ production
- ➤ Radiative decay rate can EM really be slower than weak?
- ➤ Who else could confirm? CDF / D0, Belle/Babar, BTeV /LHCb, Compass Selex (pD+K-)

Selex observes two Pair of Doubly Charmed Baryons: Two Isodoublets?





Peter S. Cooper - Fermilab